

LISTING OF PEER-REVIEWED SPMD-RELATED PUBLICATIONS:
JOURNAL ARTICLES, BOOK CHAPTERS, etc.
(as of May 2009)
by Randal Clark

n = 333

- Ahmad, N.; Bugueno, G.; Guo, L. and Marolt, R.S. 1999. Determination of organochlorine and organophosphate pesticide residues in fruits, vegetables and sediments. *Journal of Environmental Science and Health Part B-Pesticides Food Contaminants and Agricultural Wastes* 34 (5): 829-848.
- Ahn, S.; Werner, D. and Luthy, R.G. 2005. Physicochemical characterization of coke-plant soil for the assessment of polycyclic aromatic hydrocarbon availability and the feasibility of phytoremediation. *Environmental Toxicology and Chemistry* 24 (9): 2185-2195.
- Allan, I.J.; Vrana, B.; Greenwood, R.; Mills, G.A.; Roig, B. and Gonzalez, C. 2006. A "toolbox" for biological and chemical monitoring requirements for the European Union's Water Framework Directive. *Talanta* 69 (2): 302-322.
- Anderson, K.A. and Johnson, E.R. 2001. Bioavailable organochlorine pesticides in a semi-arid region of Eastern Oregon, USA, as determined by gas chromatography with electron-capture detection. *Journal of AOAC International* 84 (5): 1371-1382.
- Awata, H.; Cobb, G.P. and Anderson, T.A. 2000. A chemical test for determining biological availability of aged chemicals in soil. *International Journal of Environmental Analytical Chemistry* 78 (1): 41-49.
- Axelman, J.; Naes, K.; Naf, C. and Broman, D. 1999. Accumulation of polycyclic aromatic hydrocarbons in semipermeable membrane devices and caged mussels (*Mytilus edulis* L.) in relation to water column phase distribution. *Environmental Toxicology and Chemistry* 18 (11): 2454-2461.
- Balmer, M.E.; Poiger, T.; Droz, C.; Romanin, K.; Bergqvist, P.-A.; Müller, M.D. and Buser, H.-R. 2004. Occurrence of methyl triclosan, a transformation product of the bactericide triclosan, in fish from various lakes in Switzerland. *Environmental Science and Technology* 38 (2): 390-395.
- Balmer, M.E.; Buser, H.-R.; Müller, M.D. and Poiger, T. 2005. Occurrence of some organic UV filters in wastewater, in surface waters, and in fish from Swiss lakes. *Environmental Science and Technology* 39 (4): 953-962.
- Barber, J.L.; Sweetman, A.J.; van Wijk, D. and Jones, K.C. 2005. Hexachlorobenzene in the global environment: Emissions, levels, distribution, trends and processes. *Science of the Total Environment* 349 (1-3): 1-44.
- Barber, L.B.; Keefe, S.H.; Antweiler, R.C.; Taylor, H.E. and Wass, R.D. 2006. Accumulation of contaminants in fish from wastewater treatment wetlands. *Environmental Science and Technology* 40 (2): 603-611.
- Bartkow, M.E.; Hawker, D.W.; Kennedy, K.E. and Müller, J.F. 2004. Characterizing uptake kinetics of PAHs from the air using polyethylene-based passive air samplers of multiple surface area-to-volume ratios. *Environmental Science and Technology* 38 (9): 2701-2706.
- Bartkow, M.E.; Huckins, J.N. and Müller, J.F. 2004. Field-based evaluation of semipermeable membrane devices (SPMDs) as passive air samplers of polycyclic aromatic hydrocarbons (PAHs). *Atmospheric Environment* 38 (35): 5983-5990.
- Bartkow, M.E.; Booij, K.; Kennedy, K.E.; Müller, J.F. and Hawker, D.W. 2005. Passive air sampling theory for semivolatile organic compounds. *Chemosphere* 60 (2): 170-176.
- Bartkow, M.E.; Jones, K.C.; Kennedy, K.E.; Hawker, D.W. and Mueller, J.F. 2006. Evaluation of performance reference compounds in polyethylene-based passive air samplers. *Environmental Pollution* 144 (2): 365-370.
- Bartkow, M.E.; Kennedy, K.E.; Huckins, J.N.; Holling, N.; Komarova, T. and Müller, J.F. 2006. Photodegradation of polycyclic aromatic hydrocarbons in passive air samplers: Field testing different deployment chambers. *Environmental Pollution* 144 (2): 371-376.
- Baussant, T.; Sanni, S.; Jonsson, G.; Skadsheim, A. and Børseth, J.F. 2001. Bioaccumulation of polycyclic aromatic compounds: 1. Bioconcentration in two marine species and in semipermeable membrane devices during chronic exposure to dispersed crude oil. *Environmental Toxicology and Chemistry* 20 (6): 1175-1184.

- Bennett, E.R.; Metcalfe, T.L. and Metcalfe, C.D. 1996. Semi-permeable membrane devices (SPMDs) for monitoring organic contaminants in the Otonabee River, Ontario. *Chemosphere* 33 (3): 363-375.
- Bennett, E.R. and Metcalfe, C.D. 2000. Distribution of degradation products of alkylphenol ethoxylates near sewage treatment plants in the lower Great Lakes, North America. *Environmental Toxicology and Chemistry* 19 (4): 784-792.
- Bergknut, M.; Sehlin, E.; Lundstedt, S.; Andersson, P.L.; Haglund, P. and Tysklind, M. 2007. Comparison of techniques for estimating PAH bioavailability: Uptake in *Eisenia fetida*, passive samplers and leaching using various solvents and additives. *Environmental Pollution* 145 (1): 154-160.
- Bergqvist, P.-A.; Strandberg, B.; Hjelt, M. and Rappe, C. 1990. Nondestructive steps for the lipid reduction in PCDD and PCDF analyses. *Organohalogen Compounds* 2: 103-106.
- Bergqvist, P.-A.; Strandberg, B.; Ekelund, R.; Rappe, C. and Granmo, A. 1998. Temporal monitoring of organochlorine compounds in seawater by semipermeable membranes following a flooding episode in western Europe. *Environmental Science and Technology* 32 (24): 3887-3892.
- Bergqvist, P.-A.; Strandberg, B. and Rappe, C. 1999. Lipid removal using semipermeable membranes (SPMDs) in PCDD and PCDF analysis of fat-rich environmental samples. *Chemosphere* 38 (5): 933-943.
- Bergqvist, P.-A.; Augulyte, L. and Jurjoniene, V. 2006. PAH and PCB removal efficiencies in Umea (Sweden) and Siauliai (Lithuania) municipal wastewater treatment plants. *Water Air and Soil Pollution* 175 (1-4): 291-303.
- Blevins, D. and Fairchild, J.F. 2001. Applicability of NASQAN data for ecosystem assessments on the Missouri River. *Hydrological Processes* 15 (7): 1347-1362.
- Boehm, P.D.; Page, D.S.; Brown, J.S.; Neff, J.M. and Bence, A.E. 2005. Comparison of mussels and semi-permeable membrane devices as intertidal monitors of polycyclic aromatic hydrocarbons at oil spill sites. *Marine Pollution Bulletin* 50 (7): 740-750.
- Boehm, P.D.; Page, D.S.; Brown, J.S.; Neff, J.M. and Bence, A.E. 2006. Comparison of mussels and semi-permeable membrane devices as inter-tidal monitors of polycyclic aromatic hydrocarbons at oil spill sites (Boehm et al., 2005) - Response to Schwartz. *Marine Pollution Bulletin* 52 (1): 122-122.
- Bonetta, S., Carraro, E., Bonetta, Si, Pignata, C., Pavan, I., Romano, C. and Gilli, G. (2009) Application of semipermeable membrane device (SPMD) to assess air genotoxicity in an occupational environment. *Chemosphere*, 75 (11), 1446-1452.
- Booij, K.; Sleiderink, H.M. and Smedes, F. 1998. Calibrating the uptake kinetics of semipermeable membrane devices using exposure standards. *Environmental Toxicology and Chemistry* 17 (7): 1236-1245.
- Booij, K.; Zegers, B.N. and Boon, J.P. 2000. Levels of some polybrominated diphenyl ether (PBDE) flame retardants along the Dutch coast as derived from their accumulation in SPMDs and blue mussels (*Mytilus edulis*). *Organohalogen Compounds* 47: 89-92.
- Booij, K. and van Drooge, B.L. 2001. Polychlorinated biphenyls and hexachlorobenzene in atmosphere, sea-surface microlayer, and water measured with semi-permeable membrane devices (SPMDs). *Chemosphere* 44 (2): 91-98.
- Booij, K.; Smedes, F. and van Weerlee, E.M. 2002. Spiking of performance reference compounds in low density polyethylene and silicone passive water samplers. *Chemosphere* 46 (8): 1157-1161.
- Booij, K.; Zegers, B.N. and Boon, J.P. 2002. Levels of some polybrominated diphenyl ether (PBDE) flame retardants along the Dutch coast as derived from their accumulation in SPMDs and blue mussels (*Mytilus edulis*). *Chemosphere* 46 (5): 683-688.
- Booij, K.; Hofmans, H.E.; Fischer, C.V. and van Weerlee, E.M. 2003. Temperature-dependent uptake rates of nonpolar organic compounds by semipermeable membrane devices and low-density polyethylene membranes. *Environmental Science and Technology* 37 (2): 361-366.
- Booij, K.; Smedes, F.; van Weerlee, E.M. and Honkoop, P.J.C. 2006. Environmental monitoring of hydrophobic organic contaminants: The case of mussels versus semipermeable membrane devices. *Environmental Science and Technology* 40 (12): 3893-3900.
- Booij, K.; van Bommel, R.; Mets, A. and Dekker, R. 2006. Little effect of excessive biofouling on the uptake of organic contaminants by semipermeable membrane devices. *Chemosphere* 65 (11): 2485-2492.
- Bridges, C.M. and Little, E.E. 2003. *Using Semipermeable Membrane Devices (SPMDs) to Assess the Toxicity and Teratogenicity of Aquatic Amphibian Habitats*. In: *ASTM Special Technical*

Publication No. 1443, pp.159-168.

- Bridges, C.M.; Little, E.E.; Gardiner, D.; Petty, J.D. and Huckins, J.N. 2004. Assessing the toxicity and teratogenicity of pond water in north-central Minnesota to amphibians. *Environmental Science and Pollution Research* 11 (4): 233-239.
- Bugliosi, E. and Giorgio, G. 2000. Development of a research strategy for assessing the presence of endocrine disruptors in aquatic environment using semipermeable membrane devices (SPMDS). *Epidemiology* 11 (4): S141-S141.
- Buschini, A., Giordani, F., Pellacani, C., Rossi, C. and Poli, P. (2008) Cytotoxic and genotoxic potential of drinking water: A comparison between two different concentration methods. *Water Research*, 42 (8-9), 1999-2006.
- Čáslavský, J.; Zdrahal, Z. and Vytopilova, M. 2000. Application of SPMDs for PAH sampling in the DEZA chemical factory. *Polycyclic Aromatic Compounds* 20 (1-4): 123-141.
- Čáslavský, J. and Kotlaříková, P. 2003. High-molecular-weight polycyclic aromatic hydrocarbons in the area and vicinity of the DEZA Chemical Plant, Czech Republic. *Polycyclic Aromatic Compounds* 23 (3): 327-352.
- Čáslavský, J.; Kotlaříková, P. and Benešová, K. 2004. Sampling of airborne polycyclic aromatic hydrocarbons with semipermeable membrane devices. *Environmental Chemistry Letters* 2 (2): 89-92.
- Chambers, D.B. 1999. Semipermeable membrane devices used to estimate bioconcentration of polychlorinated biphenyls. *Journal of the American Water Resources Association* 35 (1): 143-153.
- Charlestra, L., Courtemanch, D. L., Amirbahman, A. and Patterson, H. (2008) Semipermeable membrane device (SPMD) for monitoring PCDD and PCDF levels from a paper mill effluent in the Androscoggin River, Maine, USA. *Chemosphere*, 72 (8), 1171-1180.
- Cheć, E.; Podgórska, B. and Węgrzyn, G. 2006. Direct addition of cultures of tester bacteria into semi-permeable membrane devices (SPMDs) as a modified procedure for preliminary detection of mutagenic pollution of the marine environment by use of microbiological mutagenicity assays. *Mutation Research - Genetic Toxicology and Environmental Mutagenesis* 611 (1-2): 17-24.
- Chimuka, L. and Cukrowska, E. 2006. The role of passive samplers in the monitoring of aquatic ecosystems and occupational hygiene pollution. *LC GC North America* 24 (6): 554-+.
- Chimuka, L.; Cukrowska, E. and Majors, R.E. 2006. The role of passive samplers in the monitoring of aquatic ecosystems and occupational hygiene pollution. *LC GC Europe* 19 (7): 402-+.
- Chimuka, L., Cukrowska, E. and Tutu, H. (2008) Monitoring of aquatic environments using passive samplers. *The Open Analytical Chemistry Journal*, 2 1-9.
- Cicinaite, A.; Huckins, J.N.; Alvarez, D.A.; Cranor, W.L.; Gale, R.W.; Kauneliene, V. and Bergqvist, P.-A. 2007. Feasibility of a simple laboratory approach for determining temperature influence on SPMD-air partition coefficients of selected compounds. *Atmospheric Environment*, 41 (13), 2844-2850.
- Cleveland, L.; Little, E.E.; Petty, J.D.; Johnson, B.T.; Lebo, J.A.; Orazio, C.E.; Dionne, J. and Crockett, A.B. 1997. Toxicological and chemical screening of Antarctica sediments: Use of whole sediment toxicity tests, Microtox, Mutatox and semipermeable membrane devices (SPMDs). *Marine Pollution Bulletin* 34 (3): 194-202.
- Corrigan, B.P. and Jones, K.C. 2002. Passive sampling survey of PBDEs and PCBs in UK air. *Organohalogen Compounds* 56: 449-452.
- Cranor, W. L., Alvarez, D. A., Huckins, J. N. and Petty, J. D. (2009) Uptake rate constants and partition coefficients for vapor phase organic chemicals using semipermeable membrane devices (SPMDs). *Atmospheric Environment*, 43 (20), 3211-3219.
- Crunkilton, R.L. and DeVita, W.M. 1997. Determination of aqueous concentrations of polycyclic aromatic hydrocarbons (PAHs) in an urban stream. *Chemosphere* 35 (7): 1447-1463.
- de la Torre, A.I.; Fernandez, C.; Tarazona, J.V. and Muñoz, M.J. 1995. Detection of aroclor, DDT, malathion and HCB using semipermeable membranes as concentration method. *Chemosphere* 31 (2): 2727-2737.
- de la Torre, A.I.; Fernandez, C.; Tarazona, J.V. and Muñoz, M.J. 1998. Combination of semipermeable membranes and toxicity bioassays for the detection of lipophilic toxic chemicals in environmental samples. *Ecotoxicology and Environmental Restoration*.
- de la Torre, A.I.; Jimenez, J.A.; Carballo, M.; Fernandez, C.; Roset, J. and Muñoz, M.J. 2000. Ecotoxicological evaluation of pig slurry. *Chemosphere* 41 (10): 1629-1635.

- DeVita, W.M. and Crunkilton, R.L. 1998. Quality control associated with use of semipermeable membrane devices. In: *Environmental Toxicology and Risk Assessment: Seventh Volume*. Little, E.E.; DeLonay, A.J.; Greenberg, B.M. (Eds.). American Society for Testing and Materials, West Conshocken, PA (USA), ASTM STP 1333, pp. 237-245.
- Durell, G.S.; Utvik, T.I.R.; Johnsen, S.; Frost, T. and Neff, J.M. 2006. Oil well produced water discharges to the North Sea. Part I: Comparison of deployed mussels (*Mytilus edulis*), semi-permeable membrane devices, and the DREAM model predictions to estimate the dispersion of polycyclic aromatic hydrocarbons. *Marine Environmental Research* 62 (3): 194-223.
- Echols, K.R.; Gale, R.W.; Schwartz, T.R.; Huckins, J.N.; Williams, L.L.; Meadows, J.C.; Morse, D.; Petty, J.D.; Orazio, C.E. and Tillitt, D.E. 2000. Comparing polychlorinated biphenyl concentrations and patterns in the Saginaw River using sediment, caged fish, and semipermeable membrane devices. *Environmental Science and Technology* 34 (19): 4095-4102.
- Ellis, G.S.; Huckins, J.N.; Rostad, C.E.; Schmitt, C.J.; Petty, J.D. and MacCarthy, P. 1995. Evaluation of lipid-containing semipermeable-membrane devices for monitoring organochlorine contaminants in the Upper Mississippi River. *Environmental Toxicology and Chemistry* 14 (11): 1875-1884.
- Ellis, S. G., Booij, K. and Kaputa, M. (2008) Comparison of semipermeable membrane device (SPMD) and large-volume solid-phase extraction techniques to measure water concentrations of 4,4'-DDT, 4,4'-DDE, and 4,4'-DDD in Lake Chelan, Washington. *Chemosphere*, 72 (8), 1112-1117.
- Esteve-Turrillas, F.A.; Pastor, A. and de la Guardia, M. 2006. Microwave-assisted extraction of pyrethroid insecticides from semi permeable membrane devices (SPMDs) used to indoor air monitoring. *Analytica Chimica Acta* 560 (1-2): 118-127.
- Esteve-Turrillas, F. A., Pastor, A. and de la Guardia, M. (2008) Evaluation of working air quality by using semipermeable membrane devices: Analysis of organophosphorus pesticides. *Analytica Chimica Acta*, 626 (1), 21-27. Følsvik, N.; Brevik, E.M. and Berge, J.A. 2000. Monitoring of organotin compounds in seawater using semipermeable membrane devices (SPMDs) - tentative results. *Journal of Environmental Monitoring* 2 (4): 281-284.
- Esteve-Turrillas, F. A.; Pastor, A.; and de la Guardia, M. (2007) Behaviour of semipermeable membrane devices in neutral pesticide uptake from waters. *Analytical and Bioanalytical Chemistry*, 387 (6), 2153-2162.
- Esteve-Turrillas, F. A.; Pastor, A.; and de la Guardia, M. (2007) Assessing air quality inside vehicles and at filling stations by monitoring benzene, toluene, ethylbenzene and xylenes with the use of semipermeable devices. *Analytica Chimica Acta*, 593 108-116.
- Følsvik, N.; Brevik, E.M. and Berge, J.A. 2002. Organotin compounds in a Norwegian fjord. A comparison of concentration levels in semipermeable membrane devices (SPMDs), blue mussels (*Mytilus edulis*) and water samples. *Journal of Environmental Monitoring* 4 (2): 280-283.
- Franke, S.; Meyer, C.; Heinzel, N.; Gatermann, R.; Hühnerfuss, H.; Rimkus, G.; Konig, W.A. and Francke, W. 1999. Enantiomeric composition of the polycyclic musks HHCB and AHTN in different aquatic species. *Chirality* 11 (10): 795-801.
- Gale, R.W.; Huckins, J.N.; Petty, J.D.; Peterman, P.H.; Williams, L.L.; Morse, D.; Schwartz, T.R. and Tillitt, D.E. 1997. Comparison of the uptake of dioxin like compounds by caged channel catfish and semipermeable membrane devices in the Saginaw River, Michigan. *Environmental Science and Technology* 31 (1): 178-187.
- Gale, R.W. 1998. Three-compartment model for contaminant accumulation by semipermeable membrane devices. *Environmental Science and Technology* 32 (15): 2292-2300.
- Gatermann, R.; Biselli, S.; Hühnerfuss, H.; Rimkus, G.; Franke, S.; Hecker, M.; Kallenborn, R.; Karbe, L. and König, W.A. 2002. Synthetic musks in the environment. Part 2: Enantioselective transformation of the polycyclic musk fragrances HHCB, AHTN, AHDI, and ATII in freshwater fish. *Archives of Environmental Contamination and Toxicology* 42 (4): 447-453.
- Gatermann, R.; Biselli, S.; Hühnerfuss, H.; Rimkus, G.; Hecker, M. and Karbe, L. 2002. Synthetic musks in the environment. Part 1: Species-dependent bioaccumulation of polycyclic and nitro musk fragrances in freshwater fish and mussels. *Archives of Environmental Contamination and Toxicology* 42 (4): 437-446.
- Gilli, G.; Schiliro, T.; Pignata, C.; Traversi, D.; Carraro, E.; Baiocchi, C.; Aigotti, R.; Giacosa, D. and Fea, E. 2005. Application of semipermeable membrane device for assessing toxicity in drinking water. *Chemosphere* 61 (11): 1691-1699.
- Gioia, R.; Steinnes, E.; Thomas, G.O.; Mejier, S.N. and Jones, K.C. 2006. Persistent organic pollutants in

- European background air: Derivation of temporal and latitudinal trends. *Journal of Environmental Monitoring* 8 (7): 700-710.
- Goodbred, S. L., Bryant, W. L., Rosen, M. R., Alvarez, D. A. and Spencer, T. (2009) How useful are the "other" semipermeable membrane devices (SPMDs); the mini-unit (15.2 cm long)? *Science of The Total Environment*, 407 (13), 4149-4156.
- Górecki, T. and Namieśnik, J. 2002. Passive sampling. *Trac-Trends in Analytical Chemistry* 21 (4): 276-291.
- Gouin, T.; Harner, T.; Blanchard, P.E. and MacKay, D. 2005. Passive and active air samplers as complementary methods for investigating persistent organic pollutants in the Great Lakes basin. *Environmental Science and Technology* 39 (23): 9115-9122.
- Gourlay, C.; Miege, C.; Noir, A.; Ravelet, C.; Garric, J. and Mouchel, J.-M. 2005. How accurately do semi-permeable membrane devices measure the bioavailability of polycyclic aromatic hydrocarbons to Daphnia magna? *Chemosphere* 61 (11): 1734-1739.
- Gourlay-Francé, C.; Lorgeoux, C.; and Tusseau-Vuillemin, M-H. (2008) Polycyclic aromatic hydrocarbon sampling in wastewaters using semipermeable membrane devices: Accuracy of time-weighted average concentration estimations of truly dissolved compounds. *Chemosphere*, 73 (8), 1194-1200.
- Granmo, A.; Ekelund, R.; Berggren, M.; Brorström-Lundén, E. and Bergqvist, P.-A. 2000. Temporal trend of organochlorine marine pollution indicated by concentrations in mussels, semipermeable membrane devices, and sediment. *Environmental Science and Technology* 34 (16): 3323-3329.
- Gustafson, K.E. and Dickhut, R.M. 1997. Distribution of polycyclic aromatic hydrocarbons in southern Chesapeake Bay surface water: Evaluation of three methods for determining freely dissolved water concentrations. *Environmental Toxicology and Chemistry* 16 (3): 452-461.
- Gustavson, K.E.; DeVita, W.M.; Revis, A. and Harkin, J.M. 2000. Novel use of a dual-zone restricted access sorbent: normal-phase solid-phase extraction separation of methyl oleate from polynuclear aromatic hydrocarbons stemming from semi-permeable membrane devices. *Journal of Chromatography A* 883 (1-2): 143-149.
- Gustavson, K.E. and Harkin, J.M. 2000. Comparison of sampling techniques and evaluation of semipermeable membrane devices (SPMDs) for monitoring polynuclear aromatic hydrocarbons (PAHs) in groundwater. *Environmental Science and Technology* 34 (20): 4445-4451.
- Harman, C., Tollefsen, K. E., Bøyum, O., Thomas, K. and Grung, M. (2008) Uptake rates of alkylphenols, PAHs and carbazoles in semipermeable membrane devices (SPMDs) and polar organic chemical integrative samplers (POCIS). *Chemosphere*, 72 (10), 1510-1516.
- Harman, C., Holth, T. F., Hylland, K., Thomas, K. and Grung, M. (2009) Relationship between polycyclic aromatic hydrocarbon (PAH) accumulation in semipermeable membrane devices and PAH bile metabolite levels in Atlantic cod (*Gadus morhua*). *Journal of Toxicology and Environmental Health - Part A: Current Issues*, 72 (3-4), 234-243.
- Harner, T.; Ikonomou, M.G.; Shoeib, M.; Stern, G.A. and Diamond, M.L. 2002. Passive air sampling results for polybrominated diphenyl ethers along an urban-rural transect. *Organohalogen Compounds* 57: 33-36.
- Harner, T.; Shoeib, M.; Diamond, M.L.; Stern, G.A. and Rosenberg, B. 2004. Using passive air samplers to assess urban - Rural trends for persistent organic pollutants. 1. Polychlorinated biphenyls and organochlorine pesticides. *Environmental Science and Technology* 38 (17): 4474-4483.
- Hajšlová, J. and Šetkova, L. 2004. Synthetic musks in bioindicators: Monitoring data of fish and human milk samples from the Czech Republic. In: *Synthetic Musk Fragrances in the Environment, Handbook of Environmental Chemistry: Anthropogenic Compounds* Rimkus, G.G. (Ed.). Springer Berlin, Heidelberg (Germany), Vol 3, Part 3X, pp. 151-187.
- Herve, S.; Paukku, R.; Paasivirta, J.; Heinonen, P. and Södergren, A. 1991. Uptake of organochlorines from lake water by hexane-filled dialysis membranes and by mussels. *Chemosphere* 22 (11): 997-1001.
- Herve, S.; Prest, H.F.; Heinonen, P.; Hyötyläinen, T.; Koistinen, J. and Paasivirta, J. 1995. Lipid-filled semipermeable membrane devices and mussels as samplers of organochlorine compounds in lake water. *Environmental Science and Pollution Research* 2 (1): 24-30.
- Hess, P. and Wells, D.E. 2001. Evaluation of dialysis as a technique for the removal of lipids prior to the GC determination of ortho- and non-ortho-chlorobiphenyls, using ¹⁴C-labelled congeners. *The Analyst* 126: 829-834.

- Hewitt, L.M.; Pryce, A.C.; Parrott, J.L.; Marlatt, V.; Wood, C.; Oakes, K. and van der Kraak, G.J. 2003. Accumulation of ligands for aryl hydrocarbon and sex steroid receptors in fish exposed to treated effluent from a bleached sulfite/groundwood pulp and paper mill. *Environmental Toxicology and Chemistry* 22 (12): 2890-2897.
- Hewitt, A.H.; Cope, W.G.; Kwak, T.J.; Augspurger, T.; Lazaro, P.R. and Shea, D. 2006. Influence of water quality and associated contaminants on survival and growth of the endangered Cape Fear shiner (*Notropis mekistocholas*). *Environmental Toxicology and Chemistry* 25 (9): 2288-2298.
- Hofelt, C.S. and Shea, D. 1997. Accumulation of organochlorine pesticides and PCBs by semipermeable membrane devices and *Mytilus edulis* in New Bedford Harbor. *Environmental Science and Technology* 31 (1): 154-159.
- Huckins, J.N.; Gibson, V.L.; Lebo, J.A.; Manuweera, G.K. and Gale, R.W. 1990. Insitu semipermeable polymeric samplers for monitoring hydrophobic contaminants in water. *Abstracts of Papers of the American Chemical Society* 200: 29-ENVR.
- Huckins, J.N.; Tubergen, M.W.; Lebo, J.A.; Gale, R.W. and Schwartz, T.R. 1990. Polymeric film dialysis in organic-solvent media for cleanup of organic contaminants. *Journal of the Association of Official Analytical Chemists* 73 (2): 290-293.
- Huckins, J.N.; Tubergen, M.W. and Manuweera, G.K. 1990. Semipermeable membrane devices containing model lipid: A new approach to monitoring the bioavailability of lipophilic contaminants and estimating their biocconcentration potential. *Chemosphere* 20 (5): 533-552.
- Huckins, J.N.; Manuweera, G.K.; Petty, J.D.; Lebo, J.A.; Gibson, V.L. and Meadows, J.C. 1992. Background and model development of lipid-containing semipermeable-membrane devices for estimation of time-weighted water concentrations of organic contaminants. *Abstracts of Papers of the American Chemical Society* 204: 38-ENVR.
- Huckins, J.N.; Manuweera, G.K.; Petty, J.D.; MacKay, D. and Lebo, J.A. 1993. Lipid-containing semipermeable-membrane devices for monitoring organic contaminants in water. *Environmental Science and Technology* 27 (12): 2489-2496.
- Huckins, J.N.; Petty, J.D.; Lebo, J.A.; Zajicek, J.L.; Martin, D.B.; Gibson, V.L. and Clark, R.C. 1993. Application of Passive Monitoring Technology to Multiple Environmental Media. *Abstracts of Papers of the American Chemical Society* 206: 38-ENVR.
- Huckins, J.N.; Petty, J.D.; Lebo, J.A.; Orazio, C.E.; Prest, H.F.; Tillitt, D.E.; Ellis, G.S.; Johnson, B.T. and Manuweera, G.K. 1996. Semipermeable Membrane Devices (SPMDs) for the concentration and assessment of bioavailable organic contaminants in aquatic environments. In: *Techniques in Aquatic Toxicology*. Ostrander, G.K. (Ed.). CRC Press, Boca Raton, FL (USA), pp. 625-655.
- Huckins, J.N.; Petty, J.D.; Orazio, C.E.; Lebo, J.A.; Clark, R.C.; Gibson, V.L.; Gala, W.R. and Echols, K.R. 1999. Determination of uptake kinetics (sampling rates) by lipid-containing semipermeable membrane devices (SPMDs) for polycyclic aromatic hydrocarbons (PAHs) in water. *Environmental Science and Technology* 33 (21): 3918-3923.
- Huckins, J.N.; Petty, J.D.; Lebo, J.A.; Almeida, F.V.; Booij, K.; Alvarez, D.A.; Clark, R.C. and Mogensen, B.B. 2002. Development of the permeability/performance reference compound approach for in situ calibration of semipermeable membrane devices. *Environmental Science and Technology* 36 (1): 85-91.
- Huckins, J.N.; Petty, J.D.; Prest, H.F.; Clark, R.C.; Alvarez, D.A.; Orazio, C.E.; Lebo, J.A.; Cranor, W.L. and Johnson, B.T. 2002. *A Guide for the Use of Semipermeable Membrane Devices (SPMDs) as Samplers of Waterborne Hydrophobic Organic Contaminants*. Vol. Publication No. 4690. American Petroleum Institute (API), Washington DC (USA).
- Huckins, J.N.; Petty, J.D.; Lebo, J.A.; Orazio, C.E.; Clark, R.C.; Gibson, V.L.; Chapman, D.; Huff, T. and Ganser, J. 2004. *SPMD Technology: Tutorial (3rd Edition) Online*. CERC-USGS, Columbia, MO. (USA) <http://wwwaux.cerc.cr.usgs.gov/spmd/index.htm>.
- Huckins, J.N.; Prest, H.F.; Petty, J.D.; Lebo, J.A.; Hodgins, M.M.; Clark, R.C.; Alvarez, D.A.; Gala, W.R.; Steen, A.; Gale, R.W. and Ingersoll, C.G. 2004. Overview and comparison of lipid-containing semipermeable membrane devices and oysters (*Crassostrea gigas*) for assessing organic chemical exposure. *Environmental Toxicology and Chemistry* 23 (7): 1617-1628.
- Huckins, J.N.; Petty, J.D. and Booij, K. 2006. *Monitors of Organic Chemicals in the Environment: Semipermeable Membrane Devices*. Springer Science+Business Media, New York, NY (USA) 223 pages.
- Hühnerfuss, H.; Biselli, S. and Gatermann, R. 2004. Enantioselective analysis of polycyclic musks as a

- versatile tool for the understanding of environmental processes. In: *Synthetic Musk Fragrances in the Environment, Handbook of Environmental Chemistry*. Springer-Verlag Berlin, Berlin (Germany), Vol 3, pp. 213-231.
- Hyne, R.V.; Pablo, F.; Aistropé, M.; Leonard, A.W. and Ahmad, N. 2004. Comparison of time-integrated pesticide concentrations determined from field-deployed passive samplers with daily river-water extractions. *Environmental Toxicology and Chemistry* 23 (9): 2090-2098.
- Ikonomou, M.G.; Rayne, S. and Addison, R.F. 2002. Exponential increases of the brominated flame retardants, polybrominated diphenyl ethers, in the Canadian arctic from 1981 to 2000. *Environmental Science and Technology* 36 (9): 1886-1892.
- Ikonomou, M.G.; Rayne, S.; Fischer, M.; Fernandez, M.P. and Cretney, W.J. 2002. Occurrence and congener profiles of polybrominated diphenyl ethers (PBDEs) in environmental samples from coastal British Columbia, Canada. *Chemosphere* 46 (5): 649-663.
- Isidori, M.; Fiorentino, A. and Parrella, A. 2001. Assessment of toxic potency and genotoxicity of contaminated air with SPMDs and *Tradescantia*-micronuclei assay. *Mutation Research* 483 (Suppl. 1): S167.
- Isidori, M.; Ferrara, M.; Lavorgna, M.; Nardelli, A. and Parrella, A. 2003. In situ monitoring of urban air in Southern Italy with the *Tradescantia* micronucleus bioassay and semipermeable membrane devices (SPMDs). *Chemosphere* 52 (1): 121-126.
- Jaward, F.M.; Mejier, S.N.; Steinnes, E.; Thomas, G.O. and Jones, K.C. 2004. Further studies on the latitudinal and temporal trends of persistent organic pollutants in Norwegian and UK background air. *Environmental Science and Technology* 38 (9): 2523-2530.
- Jegorova, I. 2002. Monitoring with semipermeable membrane device (SPMD) technology for persistent organic pollutants (POPs) analysis. In: *Principles of Instrumental Analysis*. Leary, J.; Skoog, D. (Eds.). Harcourt Brace Jovanovich, Orlando, FL (USA).
- Johnson, B.T.; Huckins, J.N.; Petty, J.D. and Clark, R.C. 2000. Collection and detection of lipophilic chemical contaminants in water, sediment, soil, and air - SPMD-TOX. *Environmental Toxicology* 15 (3): 248-252.
- Johnson, B.T.; Petty, J.D.; Huckins, J.N.; Lee, K. and Gauthier, J. 2004. Hazard assessment of a simulated oil spill on intertidal areas of the St. Lawrence river with SPMD-TOX. *Environmental Toxicology* 19 (4): 329-335.
- Katsoyiannis, A. and Samara, C. 2007. Comparison of active and passive sampling for the determination of persistent organic pollutants (POPs) in sewage treatment plants. *Chemosphere* 67 (7): 1375-1382.
- Kauneliene, V.; Cicenaitė, A.; Jegorova, I.; Zaliauskienė, A. and Bergqvist, P.-A. 2003. Tentative Air Concentration of PAHs and PCBs in urban area of Lithuania. *Environmental Research, Engineering, and Management* 2 (4): 33-39.
- Ke, R.H.; Li, J.; Xu, Y.P. and Wang, Z.J. 2006. Comparison of the methods for assessing Ah effects in aquatic system by semipermeable membrane device and caged fish. *Huanjing Kexue/Environmental Science* 27 (11): 2309-2313.
- Ke, R.H.; Qiao, M.; Xu, Y.P.; Huang, S.B. and Wang, Z.J. 2006. Comparison between biomimetic sampling technique using semipermeable membrane device and bioconcentration in caged fish for polycyclic aromatic hydrocarbons. *Huanjing Kexue/Environmental Science* 27 (7): 1410-1414.
- Ke, R.H.; Xu, Y.P.; Wang, Z. and Khan, S.U. 2006. Estimation of the uptake rate constants for polycyclic aromatic hydrocarbons accumulated by semipermeable membrane devices and triolein-embedded cellulose acetate membranes. *Environmental Science and Technology* 40 (12): 3906-3911.
- Ke, R.H.; Li, J.; Qiao, M.; Xu, Y.P. and Wang, Z. (2007) Using semipermeable membrane devices, bioassays, and chemical analysis for evaluation of bioavailable polycyclic aromatic hydrocarbons in water. *Archives of Environmental Contamination and Toxicology* 53 (3), 313-320.
- Kočí, V.; Ocelka, T. and Kochánková, L. 2001. SPMD - a modern approach to passive sampling. *Vodni Hospodarství* 51 (11): 331-332.
- Kočí, V.; Ocelka, T. and Kochánková, L. 2001. SPMD - Description of perspective method of POP monitoring in water bodies. *Vodni Hospodarství* 51 (12): 345-348.
- Kočí, V.; Mlejnek, M. and Kochánková, L. 2003. Toxicological evaluation of exposed SPMD membranes. *Central European Journal of Chemistry* 1 (1): 28-34.
- Kočí, V.; Lukavský, J.; Mlejnek, M.; Kochánková, L.; Grabić, R. and Ocelka, T. 2004. Application of a

- semipermeable membrane device (SPMD) for assessment of organic toxicants dangerous to the green microalga *Scenedesmus subspicatus*. *Archiv Fur Hydrobiologie - Supplement* 150: 173-186.
- Kočí, V.; Ocelka, T.; Mlejnek, M. and Grbic, R. 2004. Efficiency assessment of wastewater treatment plant based on SPMD sampling. *Central European Journal of Chemistry* 2 (1): 91-112.
- Kočí, V., Ocelka, T., Dragoun, D., Vit, M., Grbic, R. and Svab, M. (2007) Concentration of organochlorine pollutants in surface waters of the Central European biosphere reserve Krivoklatsko. *Environmental Science and Pollution Research*, 14 (2), 94-101.
- Koester, C.J.; Simonich, S.L. and Esser, B.K. 2003. Environmental analysis. *Analytical Chemistry* 75 (12): 2813-2829.
- Koester, C.J. and Moulik, A. 2005. Trends in environmental analysis. *Analytical Chemistry* 77 (12): 3737-3754.
- Koistinen, J.; Lehtonen, M.; Tukia, K.; Soimasuo, M.; Lahtipera, M. and Oikari, A. 1998. Identification of lipophilic pollutants discharged from a Finnish pulp and paper mill. *Chemosphere* 37 (2): 219-235.
- Koistinen, J.; Soimasuo, M.; Tukia, K.; Oikari, A.; Blankenship, A. and Giesy, J.P. 1998. Induction of EROD activity in HEPA-1 mouse hepatoma cells and estrogenicity in MCF-7 human breast cancer cells by extracts of pulp mill effluents, sludge, and sediment exposed to effluents. *Environmental Toxicology and Chemistry* 17 (8): 1499-1507.
- Kolok, A.S.; Huckins, J.N.; Petty, J.D. and Oris, J.T. 1996. The role of water ventilation and sediment ingestion in the uptake of benzo[a]pyrene in gizzard shad (*Dorosoma cepedianum*). *Environmental Toxicology and Chemistry* 15 (10): 1752-1759.
- Komarova, T. V.; Bartkow, M.E.; Müller, J.F.; Carter, S. and Vanderzalm, J.L. 2006. Field evaluation of passive samplers: Monitoring polycyclic aromatic hydrocarbons (PAHs) in stormwater. *Polycyclic Aromatic Compounds* 26 (3): 221-236.
- Komarova, T. V., Bartkow, M. E., Rutishauser, S., Carter, S. and Mueller, J. F. (2009) Evaluation and in situ assessment of photodegradation of polycyclic aromatic hydrocarbons in semipermeable membrane devices deployed in ocean water. *Environmental Pollution*, 157 (3), 731-736.
- Kot, A.; Zabiegala, B. and Namieśnik, J. 2000. Passive sampling for long-term monitoring of organic pollutants in water. *Trac-Trends in Analytical Chemistry* 19 (7): 446-459.
- Kukkonen, J.V.K.; Rantalainen, A.-L.; Mikkelson, P.; Lytykäinen, M.; Hämäläinen, H. and Paasivirta, J. 1998. Bioavailability of polychlorinated diphenylethers (PDPEs) in contaminated sediments: Comparison of benthic organisms and semipermeable membrane devices (SPMDs). *Organohalogen Compounds* 39: 5-8.
- Lebo, J.A.; Zajicek, J.L.; Huckins, J.N.; Petty, J.D. and Peterman, P.H. 1992. Use of semipermeable-membrane devices for insitu monitoring of polycyclic aromatic-hydrocarbons in aquatic environments. *Chemosphere* 25 (5): 697-718.
- Lebo, J.A.; Gale, R.W.; Petty, J.D.; Tillitt, D.E.; Huckins, J.N.; Meadows, J.C.; Echols, K.R. and Immon, L.E. 1994. Analyses of PCDD PCDF residues sequestered from environmental waters by the semipermeable-membrane device. *Abstracts of Papers of the American Chemical Society* 208: 226.
- Lebo, J.A.; Gale, R.W.; Petty, J.D.; Tillitt, D.E.; Huckins, J.N.; Meadows, J.C.; Orazio, C.E.; Echols, K.R.; Schroeder, D.J. and Immon, L.E. 1995. Use of the semipermeable-membrane device as an in-situ sampler of waterborne bioavailable PCDD and PCDF residues at sub-parts-per-quadrillion concentrations. *Environmental Science and Technology* 29 (11): 2886-2892.
- Lebo, J.A.; Zajicek, J.L.; Orazio, C.E.; Petty, J.D.; Huckins, J.N. and Douglas, E.H. 1996. Use of the semipermeable membrane device (SPMD) to sample polycyclic aromatic hydrocarbon pollution in a lotic system. *Polycyclic Aromatic Compounds* 8 (1): 53-65.
- Lebo, J.A.; Huckins, J.N.; Petty, J.D. and Ho, K.T. 1999. Removal of organic contaminant toxicity from sediments - Early work toward development of a toxicity identification evaluation (TIE) method. *Chemosphere* 39 (3): 389-406.
- Lebo, J.A.; Huckins, J.N.; Petty, J.D.; Ho, K.T. and Stern, E.A. 2000. Selective removal of organic contaminants from sediments: A methodology for toxicity identification evaluations (TIEs). *Chemosphere* 40 (8): 811-819.
- Lebo, J.A.; Almeida, F.V.; Cranor, W.L.; Petty, J.D.; Huckins, J.N.; Rastall, A.C.; Alvarez, D.A.; Mogensen, B.B. and Johnson, B.T. 2004. Purification of triolein for use in semipermeable membrane devices (SPMDs). *Chemosphere* 54 (8): 1217-1224.

- Leiker, T. J., Abney, S. R., Goodbred, S. L. and Rosen, M. R. (2009) Identification of methyl triclosan and halogenated analogues in male common carp (*Cyprinus carpio*) from Las Vegas Bay and semipermeable membrane devices from Las Vegas Wash, Nevada. *Science of The Total Environment*, 407 (6), 2102-2114.
- Leppänen, M.T. and Kukkonen, J.V.K. 2000. Effect of sediment-chemical contact time on availability of sediment-associated pyrene and benzo[a]pyrene to oligochaete worms and semi-permeable membrane devices. *Aquatic Toxicology* 49 (4): 227-241.
- Leppänen, M.T. and Kukkonen, J.V.K. 2006. Evaluating the role of desorption in bioavailability of sediment-associated contaminants using oligochaetes, semipermeable membrane devices and Tenax extraction. *Environmental Pollution* 140 (1): 150-163.
- Li, J.; Zhang, G.; Qi, S. and Liu, G.Q. 2004. Exchange of polycyclic aromatic hydrocarbons (PAHs) across air-water interface in Luhu Lake, Guangzhou. *Hupo Kexue* 16 (3): 238-244.
- Liao, L.B. and Xiao, X.M. 2006. Accumulation of organochlorine pesticides by semipermeable membrane devices using composite complex. *Chemosphere* 64 (9): 1592-1600.
- Lico, M.S. 2004. Gasoline-related organics in Lake Tahoe before and after prohibition of carbureted two-stroke engines. *Lake and Reservoir Management* 20 (2): 164-174.
- Lindström, A.; Buerge, I.J.; Poiger, T.; Bergqvist, P.-A.; Müller, M.D. and Buser, H.-R. 2002. Occurrence and environmental behavior of the bactericide triclosan and its methyl derivative in surface waters and in wastewater. *Environmental Science and Technology* 36 (11): 2322-2329.
- Liu, G.Q.; Zhang, G.; Li, J.; Peng, X. and Qi, S. 2004. Studies on monitoring atmospheric PAHs utilizing SPMD technique in the Pearl River Delta. *China Environmental Science* 24 (3): 340-344.
- Liu, G.Q.; Zhang, G.; Li, J.; Peng, X. and Qi, S. 2004. Monitoring OCPs in the air using SPMD in the Pearl River delta. *Research on environmental science/Huanjing Kexue Yanjiu* 17 (6): 1-4,11.
- Liu, G.Q.; Zhang, G.; Li, J.; Li, X.; Peng, X. and Qi, S. 2006. Spatial distribution and seasonal variations of polycyclic aromatic hydrocarbons (PAHs) using semi-permeable membrane devices (SPMD) and pine needles in the Pearl River Delta, South China. *Atmospheric Environment* 40 (17): 3134-3143.
- Lohmann, R.; Corrigan, B.P.; Howsam, M.; Jones, K.C. and Ockenden, W.A. 2001. Further developments in the use of semipermeable membrane devices (SPMDs) as passive air samplers for persistent organic pollutants: Field application in a spatial survey of PCDD/Fs and PAHs. *Environmental Science and Technology* 35 (12): 2576-2582.
- Louch, J.; Allen, G.; Erickson, C.; Wilson, G. and Schmedding, D. 2003. Interpreting results from field deployments of semipermeable membrane devices. *Environmental Science and Technology* 37 (6): 1202-1207.
- Lu, Y. and Wang, Z. 2002. Bioconcentration of trace organochlorine pesticides by the rainbow trout. *Journal of Environmental Science and Health Part a-Toxic/Hazardous Substances & Environmental Engineering* 37 (4): 529-539.
- Lu, Y.; Wang, Z. and Huckins, J.N. 2002. Review of the background and application of triolein-containing semipermeable membrane devices in aquatic environmental study. *Aquatic Toxicology* 60 (1-2): 139-153.
- Lu, Y. and Wang, Z. 2003. Accumulation of organochlorinated pesticides by triolein-containing semipermeable membrane device (triolein-SPMD) and rainbow trout. *Water Research* 37 (10): 2419-2425.
- Lu, Y. and Wang, Z. 2004. Accumulation kinetics of organochlorinated pesticides by triolein-containing semipermeable membrane devices. *Acta Hydrobiologica Sinica* 28 (5): 540-544.
- Luellen, D.R. and Shea, D. 2002. Calibration and field verification of semipermeable membrane devices for measuring polycyclic aromatic hydrocarbons in water. *Environmental Science and Technology* 36 (8): 1791-1797.
- Luellen, D.R. and Shea, D. 2003. Semipermeable membrane devices accumulate conserved ratios of sterane and hopane petroleum biomarkers. *Chemosphere* 53 (7): 705-713.
- Lyttikäinen, M.; Hirva, P.; Minkkinen, P.; Hämäläinen, H.; Rantalainen, A.-L.; Mikkelson, P.; Paasivirta, J. and Kukkonen, J.V.K. 2003. Bioavailability of sediment-associated PCDD/Fs and PCDEs: Relative importance of contaminant and sediment characteristics and biological factors. *Environmental Science and Technology* 37 (17): 3926-3934.
- Lyttikäinen, M.; Rantalainen, A.-L.; Mikkelson, P.; Hämäläinen, H.; Paasivirta, J. and Kukkonen, J.V.K. 2003. Similarities in bioaccumulation patterns of polychlorinated dibenz-p-dioxins and furans

- and polychlorinated diphenyl ethers in laboratory-exposed oligochaetes and semipermeable membrane devices and in field-collected chironomids. *Environmental Toxicology and Chemistry* 22 (10): 2405-2415.
- Ma, M.; Wang, Y. and Wang, Z. 1999. Sample pretreatment of toxic assessment of organic pollutants in river water. *Huanjing Kexue/Environmental Science* 20 (6): 80-83.
- Ma, M. and Wang, Z. 2004. Assessing toxicity of organic components in water and wastewater using active and passive sampling and luminescent bacteria bioassays. *Acta Scientiae Circumstantiae/Huanjing Xuebao* 24 (4): 684-689.
- Ma, M.; Wang, C. and Wang, Z. 2005. Assessing toxicities of hydrophobic organic pollutants in Huaihe River by using two types of sampling. *Journal of Environmental Science and Health Part a-Toxic/Hazardous Substances & Environmental Engineering* 40 (2): 331-342.
- MacKay, D. 2004. Finding fugacity feasible, fruitful, and fun. *Environmental Toxicology and Chemistry* 23 (10): 2282-2289.
- MacRae, J.D. and Hall, K.J. 1998. Comparison of methods used to determine the availability of polycyclic aromatic hydrocarbons in marine sediment. *Environmental Science and Technology* 32 (23): 3809-3815.
- McCarthy, J.F.; Southworth, G.R.; Ham, K.D. and Palmer, J.A. 2000. Time-integrated, flux-based monitoring using semipermeable membrane devices to estimate the contribution of industrial facilities to regional polychlorinated biphenyl budgets. *Environmental Toxicology and Chemistry* 19 (2): 352-359.
- McCarthy, K.A. and Gale, R.W. 2001. Evaluation of persistent hydrophobic organic compounds in the Columbia River Basin using semipermeable-membrane devices. *Hydrological Processes* 15 (7): 1271-1283.
- McCarthy, K. 2006. Assessment of the usefulness of semipermeable membrane devices for long-term watershed monitoring in an urban slough system. *Environmental Monitoring and Assessment* 118 (1-3): 293-318.
- Meadows, J.C.; Tillitt, D.E.; Huckins, J.N. and Schroeder, D.J. 1993. Large-scale dialysis of sample lipids using a semipermeable-membrane device. *Chemosphere* 26 (11): 1993-2006.
- Meadows, J.C.; Tillitt, D.E.; Schwartz, T.R.; Schroeder, D.J.; Echols, K.R.; Gale, R.W.; Powell, D.C. and Bursian, S.J. 1996. Organochlorine contaminants in double-crested cormorants from Green Bay, WI. I. Large-scale extraction and isolation from eggs using semi-permeable membrane dialysis. *Archives of Environmental Contamination and Toxicology* 31 (2): 218-224.
- Meadows, J.C.; Echols, K.R.; Huckins, J.N.; Borsuk, F.A.; Carline, R.F. and Tillitt, D.E. 1998. Estimation of uptake rate constants for PCB congeners accumulated by semipermeable membrane devices and brown trout (*Salmo trutta*). *Environmental Science and Technology* 32 (12): 1847-1852.
- Meijer, S.N.; Ockenden, W.A.; Steinnes, E.; Corrigan, B.P. and Jones, K.C. 2003. Spatial and temporal trends of POPs in Norwegian and UK background air: Implications for global cycling. *Environmental Science and Technology* 37 (3): 454-461.
- Metcalfe, T.L.; Metcalfe, C.D.; Bennett, E.R. and Haffner, G.D. 2000. Distribution of toxic organic contaminants in water and sediments in the Detroit river. *Journal of Great Lakes Research* 26 (1): 55-64.
- Metcalfe, T.L.; Dillon, P.J. and Metcalfe, C.D. (2008) Detecting the transport of toxic pesticides from golf courses into watersheds in the Precambrian Shield region of Ontario, Canada. *Environmental Toxicology and Chemistry*, 27 (4), 811-812.
- Miege, C.; Durand, S.; Garric, J.; Gourlay, C.; Wang, D.; Mouchel, J.-M. and Tusseau-Vuillemin, M.H. 2004. Semipermeable membrane device-availability of polycyclic aromatic hydrocarbons in river waters and wastewater treatment plant effluents. *Polycyclic Aromatic Compounds* 24 (4-5): 805-825.
- Miege, C.; Ravelet, C.; Croue, J.P. and Garric, J. 2005. Semi-permeable membrane device efficiency for sampling free soluble fraction of polycyclic aromatic hydrocarbons. *Analytica Chimica Acta* 536 (1-2): 259-266.
- Moore, M.T.; Bennett, E.R.; Cooper, C.M.; Smith, S.; Farris, J.L.; Drouillard, K.G. and Schulz, R. 2006. Influence of vegetation in mitigation of methyl parathion runoff. *Environmental Pollution* 142 (2): 288-294.
- Moring, J.B. and Rose, D.R. 1997. Occurrence and concentrations of polycyclic aromatic hydrocarbons in semipermeable membrane devices and clams in three urban streams of the Dallas-Fort Worth

- Metropolitan Area, Texas. *Chemosphere* 34 (3): 551-566.
- Namieśnik, J.; Zabiegala, B.; Kot-Wasik, A.; Partyka, M. and Wasik, A. 2005. Passive sampling and/or extraction techniques in environmental analysis: A review. *Analytical and Bioanalytical Chemistry* 381 (2): 279-301.
- Neal, J.W.; Bachelier, N.M.; Noble, R.L.; Shea, D. and Cope, W.G. 2005. The mystery of Dos Bocas Reservoir, Puerto Rico: Explaining extreme spatial heterogeneity in largemouth bass distribution. *Caribbean Journal of Science* 41 (4): 804-814.
- Norrgren, L.; Pettersson, U.; Örn, S. and Bergqvist, P.-A. 2000. Environmental monitoring of the Kafue River, located in the Copperbelt, Zambia. *Archives of Environmental Contamination and Toxicology* 38 (3): 334-341.
- Ocelka, T.; Pazak, J.; Kočí, V.; Raclavska, H. and Grbic, R. 2003. Application SPMD method for POPs monitoring and toxicity assessment in waste waters. *Vodni Hospordarstvi* 53 (8): 211-214.
- Ocelka, T.; Kočí, V.; Grbic, R.; Hapala, P.; Raclavska, H.; Prazak, J. and Leontovycova, D. 2004. Semipermeable membrane device (SPMD) as a tool for POP's and toxicity assessment. Application I: Sewage sludge assessment and biodegradation - Application II: Monitoring of rivers in the Czech Republic. *International Biodeterioration & Biodegradation* 53 (4): 241.
- Ockenden, W.A.; Prest, H.F.; Thomas, G.O.; Sweetman, A.J. and Jones, K.C. 1998. Passive air sampling of PCBs: Field calculation of atmospheric sampling rates by triolein-containing semipermeable membrane devices. *Environmental Science and Technology* 32 (10): 1538-1543.
- Ockenden, W.A.; Sweetman, A.J.; Prest, H.F.; Steinnes, E. and Jones, K.C. 1998. Toward an understanding of the global atmospheric distribution of persistent organic pollutants: The use of semipermeable membrane devices as time-integrated passive samplers. *Environmental Science and Technology* 32 (18): 2795-2803.
- Ockenden, W.A.; Corrigan, B.P.; Howsam, M. and Jones, K.C. 2001. Further developments in the use of semipermeable membrane devices as passive air samplers: Application to PCBs. *Environmental Science and Technology* 35 (22): 4536-4543.
- O'Toole, S.; Metcalfe, C.D.; Craine, I. and Gross, M. 2006. Release of persistent organic contaminants from carcasses of Lake Ontario Chinook salmon (*Oncorhynchus tshawytscha*). *Environmental Pollution* 140 (1): 102-113.
- Pablos, M.; Fernandez, C.; Garcia-Hortiguela, P.; Valdovinos, C. and Tarazona, J.V. 1999. Comparison of different extraction procedures for organic-fraction toxicity testing of urban sewages. *Toxicological and Environmental Chemistry* 70 (1-2): 115-127.
- Palowitch, A.W. 1996. Semipermeable membrane devices. *Sea Technology* 37 (2): 73-79.
- Parrott, J.L. and Tillitt, D.E. 1997. The use of semipermeable membrane devices (SPMDs) to concentrate inducers of fish hepatic mixed function oxygenase (MFO). In: *Ecotoxicology: Responses, Biomarkers, and Risk Assessment. An OECD Workshop*. Zelikoff, J.T. (Ed.). SOS Publications, Fair Haven, NJ (USA), pp. 185-196.
- Parrott, J.L.; Backus, S.M.; Borgmann, A.I. and Swyripa, M. 1999. The use of semipermeable membrane devices to concentrate chemicals in oil refinery effluent on the Mackenzie River. *Arctic* 52 (2): 125-138.
- Paschke, A.; Vrana, B.; Popp, P. and Schuurmann, G. 2006. Comparative application of solid-phase microextraction fibre assemblies and semi-permeable membrane devices as passive air samplers for semi-volatile chlorinated organic compounds. A case study on the landfill "Grube Antonie" in Bitterfeld, Germany. *Environmental Pollution* 144 (2): 414-422.
- Petty, J.D.; Huckins, J.N. and Zajicek, J.L. 1993. Application of semipermeable membrane devices (SPMDs) as passive air samplers. *Chemosphere* 27 (9): 1609-1624.
- Petty, J.D.; Huckins, J.N.; Martin, D.B. and Adornato, T.G. 1995. Use of Semipermeable-Membrane Devices (SPMDs) to determine bioavailable organochlorine pesticide-residues in streams receiving irrigation drain water. *Chemosphere* 30 (10): 1891-1903.
- Petty, J.D.; Huckins, J.N.; Orazio, C.E.; Lebo, J.A.; Poulton, B.C.; Gale, R.W.; Charbonneau, C.S. and Kaiser, E.M. 1995. Determination of waterborne bioavailable organochlorine pesticide residues in the lower Missouri River. *Environmental Science and Technology* 29 (10): 2561-2566.
- Petty, J.D.; Poulton, B.C.; Charbonneau, C.S.; Huckins, J.N.; Jones, S.B.; Cameron, J.T. and Prest, H.F. 1998. Determination of bioavailable contaminants in the lower Missouri River following the flood of 1993. *Environmental Science and Technology* 32 (7): 837-842.
- Petty, J.D.; Jones, S.B.; Huckins, J.N.; Cranor, W.L.; Parris, J.T.; McTague, T.B. and Boyle, T.P. 2000.

- An approach for assessment of water quality using semipermeable membrane devices (SPMDs) and bioindicator tests. *Chemosphere* 41 (3): 311-321.
- Petty, J.D.; Orazio, C.E.; Huckins, J.N.; Gale, R.W.; Lebo, J.A.; Meadows, J.C.; Echols, K.R. and Cranor, W.L. 2000. Considerations involved with the use of semipermeable membrane devices for monitoring environmental contaminants. *Journal of Chromatography A* 879 (1): 83-95.
- Petty, J.D.; Huckins, J.N.; Alvarez, D.A.; Brumbaugh, W.G.; Cranor, W.L.; Gale, R.W.; Rastall, A.C.; Jones-Lepp, T.L.; Leiker, T.J.; Rostad, C.E. and Furlong, E.T. 2004. A holistic passive integrative sampling approach for assessing the presence and potential impacts of waterborne environmental contaminants. *Chemosphere* 54 (6): 695-705.
- Peven, C.S.; Uhler, A.D. and Querzoli, F.J. 1996. Caged mussels and semipermeable membrane devices as indicators of organic contaminant uptake in Dorchester and Duxbury Bays, Massachusetts. *Environmental Toxicology and Chemistry* 15 (2): 144-149.
- Poiger, T.; Buser, H.-R.; Balmer, M.E.; Bergqvist, P.-A. and Müller, M.D. 2004. Occurrence of UV filter compounds from sunscreens in surface waters: Regional mass balance in two Swiss lakes. *Chemosphere* 55 (7): 951-963.
- Prest, H.F.; Jarman, W.M.; Burns, S.A.; Weismuller, T.; Martin, M. and Huckins, J.N. 1992. Passive water sampling via semipermeable membrane device (SPMDS) in concert with bivalves in the Sacramento/San Joaquin River delta. *Chemosphere* 25 (12): 1811-1823.
- Prest, H.F.; Huckins, J.N.; Petty, J.D.; Herve, S.; Paasivirta, J. and Heinonen, P. 1995. A survey of recent results in passive sampling of water and air by semipermeable membrane devices. *Marine Pollution Bulletin* 31 (4-12): 306-312.
- Prest, H.F.; Jacobson, L.A. and Huckins, J.N. 1995. Passive sampling of water and coastal air via semipermeable-membrane devices. *Chemosphere* 30 (7): 1351-1361.
- Prest, H.F.; Richardson, B.J.; Jacobson, L.A.; Vedder, J. and Martin, M. 1995. Monitoring organochlorines with semi-permeable membrane devices (SPMDs) and mussels (*Mytilus edulis*) in Corio Bay, Victoria, Australia. *Marine Pollution Bulletin* 30 (8): 543-554.
- Prest, H.F.; Jacobson, L.A. and Wilson, M. 1997. Passive water sampling for polynuclear aromatic hydrocarbons using lipid-containing semipermeable membrane devices (SPMDs): Application to contaminant residence times. *Chemosphere* 35 (12): 3047-3063.
- Pulkarabová, J., Suchanová, M., Hajšlová, J., Kocourek, V. and Tomaniová, M. (2009) Passive sampling of water and Porous Media for monitoring of organic pollutants. *Chemicke Listy*, 103 (2), 159-165.
- Rantalainen, A.-L.; Ikonomou, M.G. and Rogers, I.H. 1998. Lipid-containing semipermeable membrane devices (SPMDs) as concentrators of toxic chemicals in the Lower Fraser River, Vancouver, British Columbia. *Chemosphere* 37 (6): 1119-1138.
- Rantalainen, A.-L.; Paasivirta, J. and Herve, S. 1998. Uptake of chlorohydrocarbons from soil by lipid-containing semipermeable membrane devices (SPMDs). *Chemosphere* 36 (6): 1415-1427.
- Rantalainen, A.-L.; Hyötyläinen, T.; Saramo, M. and Niskanen, I. 1999. Passive sampling of PAHs in indoor air in Nepal. *Toxicological and Environmental Chemistry* 68 (3-4): 335-348.
- Rantalainen, A.-L.; Cretney, W.J. and Ikonomou, M.G. 2000. Uptake rates of semipermeable membrane devices (SPMDs) for PCDDs, PCDFs and PCBs in water and sediment. *Chemosphere* 40 (2): 147-158.
- Rantalainen, A.-L.; Cretney, W.J. and Ikonomou, M.G. 2000. Comparison of three techniques for lipid removal from seal blubber: Gel permeation, acid treatment, and dialysis with semipermeable membrane. *International Journal of Environmental Analytical Chemistry* 76: 31-47.
- Rappe, C. and Granmo, A. 1998. Temporal monitoring of organochlorine compounds in seawater by semipermeable membranes following a flooding episode in Western Europe. *Environmental Science and Technology* 32 (24): 3887-3892.
- Rastall, A.C.; Neziri, A.; Vukovic, Z.; Jung, C.; Mijovic, S.; Hollert, H.; Nikcevic, S. and Erdinger, L. 2004. The identification of readily bioavailable pollutants in Lake Shkodra/Skadar using semipermeable membrane devices (SPMDs), bioassays and chemical analysis. *Environmental Science and Pollution Research* 11 (4): 240-253.
- Rastall, A.C.; Getting, D.; Goddard, J.; Roberts, D.R. and Erdinger, L. 2006. A biomimetic approach to the detection and identification of estrogen receptor agonists in surface waters using semipermeable membrane devices (SPMDs) and bioassay-directed chemical analysis. *Environmental Science and Pollution Research* 13 (4): 256-267.
- Rayne, S. and Ikonomou, M.G. 2002. Reconstructing source polybrominated diphenyl ether congener

- patterns from semipermeable membrane devices in the Fraser River, British Columbia, Canada: Comparison to commercial mixtures. *Environmental Toxicology and Chemistry* 21 (11): 2292-2300.
- Richardson, B.J.; Zheng, G.J.; Tse, E.S.C. and Lam, P.K.S. 2001. A comparison of mussels (*Perna viridis*) and semi-permeable membrane devices (SPMDs) for monitoring chlorinated trace organic contaminants in Hong Kong coastal waters. *Chemosphere* 45 (8): 1201-1208.
- Richardson, B.J.; Lam, P.K.S.; Zheng, G.J.; McClellan, K.E. and De Luca-Abbott, S.B. 2002. Biofouling confounds the uptake of trace organic contaminants by semi-permeable membrane devices (SPMDs). *Marine Pollution Bulletin* 44 (12): 1372-1379.
- Richardson, B.J.; Zheng, G.J.; Tse, E.S.C.; De Luca-Abbott, S.B.; Siu, S.Y.M. and Lam, P.K.S. 2003. A comparison of polycyclic aromatic hydrocarbon and petroleum hydrocarbon uptake by mussels (*Perna viridis*) and semi-permeable membrane devices (SPMDs) in Hong Kong coastal waters. *Environmental Pollution* 122 (2): 223-227.
- Richardson, B.J.; Tse, E.S.C.; De Luca-Abbott, S.B.; Martin, M. and Lam, P.K.S. 2005. Uptake and depuration of PAHs and chlorinated pesticides by semi-permeable membrane devices (SPMDs) and green-lipped mussels (*Perna viridis*). *Marine Pollution Bulletin* 51 (8-12): 975-993.
- Rogers, H.R. 1997. Influence of suspended solids and back diffusion on organic contaminant uptake by semi-permeable membranes (SPMDs). *Chemosphere* 35 (8): 1651-1657.
- Rogers, K.R.; Harper, S.L. and Robertson, G.L. 2005. Screening for toxic industrial chemicals using semipermeable membrane devices with rapid toxicity assays. *Analytica Chimica Acta* 543 (1-2): 229-235.
- Rohr, A.C.; Hall, E.R. and Hall, K.J. 1996. Use of semipermeable membrane devices for monitoring pulp mill effluents: A preliminary assessment. *Water Quality Research Journal of Canada* 31 (1): 85-100.
- Rosen, M.R.; Rowe, T.G.; Goodbred, S.L.; Shipley, D.O. and Arufe, J.A. 2006. Importance of land use, streamflow, and water quality on stream toxicity in the lake tahoe and truckee river watersheds. *American Fisheries Society Symposium* 2006 (48): 129-149.
- Rostad, C.E.; Ellis, G.S.; Schmitt, C.J.; Huckins, J.N. and Petty, J.D. 1993. Insitu application of semipermeable-membrane devices for monitoring lipophilic contaminants in the upper Mississippi River. *Abstracts of Papers of the American Chemical Society* 205: 83-ENV.
- Sabaliūnas, D. and Södergren, A. 1996. Uptake of organochlorine pesticides by solvent-filled cellulose and polyethylene membranes. *Ecotoxicology and Environmental Safety* 35 (2): 150-155.
- Sabaliūnas, D. and Södergren, A. 1997. Use of semi-permeable membrane devices to monitor pollutants in water and assess their effects: A laboratory test and field verification. *Environmental Pollution* 96 (2): 195-205.
- Sabaliūnas, D.; Ellington, J. and Neuman, J. 1998. Semipermeable membrane devices for monitoring pollutants and their effects in aquatic ecosystems of Lithuania. *Critical Reviews in Analytical Chemistry* 28 (2): 50-50.
- Sabaliūnas, D.; Lazutka, J.R.; Sabaliūnienė, I. and Södergren, A. 1998. Use of semipermeable membrane devices for studying effects of organic pollutants: Comparison of pesticide uptake by semipermeable membrane devices and mussels. *Environmental Toxicology and Chemistry* 17 (9): 1815-1824.
- Sabaliūnas, D.; Ellington, J. and Sabaliūnienė, I. 1999. Screening bioavailable hydrophobic toxicants in surface waters with semipermeable membrane devices: Role of inherent oleic acid in toxicity evaluations. *Ecotoxicology and Environmental Safety* 44 (2): 160-167.
- Sabaliūnas, D.; Lazutka, J.R. and Sabaliūnienė, I. 2000. Acute toxicity and genotoxicity of aquatic hydrophobic pollutants sampled with semipermeable membrane devices. *Environmental Pollution* 109 (2): 251-265.
- Sabaliūnas, D.; Webb, S.F.; Hauk, A.; Jacob, M. and Eckhoff, W.S. 2003. Environmental fate of Triclosan in the River Aire Basin, UK. *Water Research* 37 (13): 3145-3154.
- Sanchez-Hernandez, J.C.; Borghini, F.; Corral, A. and Grimalt, J.O. 2004. Field uptake rates of hydrophobic organic contaminants by semipermeable membrane devices: Environmental monitoring considerations. *Journal of Environmental Monitoring* 6 (11): 919-925.
- Schaanning, M.; Breyholtz, B. and Skei, J. 2006. Experimental results on effects of capping on fluxes of persistent organic pollutants (POPs) from historically contaminated sediments. *Marine Chemistry* 102 (1-2): 46-59.

- Schibetta, R.C. and Manock, J.J. 2000. Use of semipermeable membrane devices (SPMDs) for the calibration of tributyltin uptake rates in water. *Abstracts of Papers of the American Chemical Society* 219: U642-U642.
- Schwartz, R.M. (2006) Comparison of mussels and semi-permeable membrane devices as inter-tidal monitors of polycyclic aromatic hydrocarbons at oil spill sites (Boehm et al. 2005) letter. *Marine Pollution Bulletin*, 52 (1), 121-121.
- Scott, G.I.; Fulton, M.H.; Wirth, E.F.; Chandler, G.T.; Key, P.B.; Daugomah, J.W.; Bearden, D.; Chung, K.W.; Strozier, E.D.; DeLorenzo, M.; Sivertsen, S.; Dias, A.; Sanders, M.; MaCauley, J.M.; Goodman, L.R.; LaCroix, M.W.; Thayer, G.W. and Kucklick, J. 2002. Toxicological studies in tropical ecosystems: An ecotoxicological risk assessment of pesticide runoff in South Florida estuarine ecosystems. *Journal of Agricultural and Food Chemistry* 50 (15): 4400-4408.
- Sethajintanin, D. and Anderson, K.A. 2006. Temporal bioavailability of organochlorine pesticides and PCBs. *Environmental Science and Technology* 40 (12): 3689-3695.
- Šetkova, L.; Hajšlová, J.; Bergqvist, P.-A.; Kocourek, V.; Kazda, R. and Suchan, P. 2005. Fast isolation of hydrophobic organic environmental contaminants from exposed semipermeable membrane devices (SPMDs) prior to GC analysis. *Journal of Chromatography A* 1092 (2): 170-181.
- Shaw, M.; Tibbetts, I.R. and Müller, J.F. 2004. Monitoring PAHs in the Brisbane River and Moreton Bay, Australia, using semipermeable membrane devices and EROD activity in yellowfin bream, *Acanthopagrus australis*. *Chemosphere* 56 (3): 237-246.
- Shaw, M. and Müller, J.F. 2005. Preliminary evaluation of the occurrence of herbicides and PAHs in the Wet Tropics region of the Great Barrier Reef, Australia, using passive samplers. *Marine Pollution Bulletin* 51 (8-12): 876-881.
- Shen, H., Henkelmann, B., Levy, W., Zsolnay, A., Weiss, P., Jakobi, G., Kirchner, M., Moche, W., Braun, K. and Schramm, K. W. (2009) Altitudinal and chiral signature of persistent organochlorine pesticides in air, soil, and spruce needles (*Picea abies*) of the alps. *Environmental Science and Technology*, 43 (7), 2450-2455.
- Shigenaka, G. and Henry, C.B., Jr. 1995. Use of mussels and semipermeable membrane devices to assess bioavailability of residual polynuclear aromatic hydrocarbons three years after the Exxon Valdez oil spill. In: *Exxon Valdez Oil Spill: Fate and Effects in Alaskan Waters*. Wells, P.G.; Butler, J.N.; Hughes, J.S. (Eds.). ASTM Philadelphia, PA (USA), ASTM STP 1219, pp. 239-260.
- Shim, S.M.; Santerre, C.R.; Dorworth, L.E.; Miller, B.K.; Stahl, J.R. and Deardorff, D.C. 2004. Prediction of PCB content in sportfish using semipermeable membrane devices (SPMDs). *Journal of Environmental Science and Health Part B-Pesticides Food Contaminants and Agricultural Wastes* 39 (2): 263-271.
- Shoeib, M. and Harner, T. 2002. Characterization and comparison of three passive air samplers for persistent organic pollutants. *Environmental Science and Technology* 36 (19): 4142-4151.
- Short, J.W.; Springman, K.R.; Lindeberg, M.R.; Holland, L.G.; Larsen, M.L.; Sloan, C.A.; Khan, C.; Hodson, P.V.; and Rice, S.D. (2008) Semipermeable membrane devices link site-specific contaminants to effects: PART II - A comparison of lingering Exxon Valdez oil with other potential sources of CYP1A inducers in Prince William Sound, Alaska. *Marine Environmental Research*, 66 (5), 487-498.
- Söderström, H.S. and Bergqvist, P.-A. 2003. Polycyclic aromatic hydrocarbons in a semiaquatic plant and semipermeable membrane devices exposed to air in Thailand. *Environmental Science and Technology* 37 (1): 47-52.
- Söderström, H.S. and Bergqvist, P.-A. 2004. Passive air sampling using semipermeable membrane devices at different wind-speeds in situ calibrated by performance reference compounds. *Environmental Science and Technology* 38 (18): 4828-4834.
- Söderström, H.S. and Bergqvist, P.-A. 2005. Wind effects on passive air sampling of PAHs and PCBs. *Bulletin of Environmental Contamination and Toxicology* 74 (2): 429-436.
- Söderström, H.S.; Hajšlová, J.; Kocourek, V.; Siegmund, B.; Kocan, A.; Obiedzinski, W.; Tysklind, M. and Bergqvist, P.-A. 2005. PAHs and nitrated PAHs in air of five European countries determined using SPMDs as passive samplers. *Atmospheric Environment* 39 (9): 1627-1640.
- Springman, K.R.; Short, J.W.; Lindeberg, M. and Rice, S.D. 2006. Induction of CYP1A in rainbow trout from bioavailable Exxon Valdez oil: Fifteen years and still counting. *Marine Environmental Research* 62 (SUPPL. 1).
- Springman, K.R.; Short, J.W.; Lindeberg, M.; and Rice, S.D. (2008) Evaluation of bioavailable

- hydrocarbon sources and their induction potential in Prince William Sound, Alaska. *Marine Environmental Research*, 66 (1), 218-220.
- Springman, K.R.; Short, J.W.; Lindeberg, M.R.; Maselko, J.M.; Khan, C.; Hodson, P.V.; and Rice, S.D. (2008) Semipermeable membrane devices link site-specific contaminants to effects: Part 1 - Induction of CYP1A in rainbow trout from contaminants in Prince William Sound, Alaska. *Marine Environmental Research*, 66 (5), 477-486.
- Strandberg, B.; Wagman, N.; Bergqvist, P.-A.; Haglund, P. and Rappe, C. 1997. Semipermeable membrane devices as passive samplers to determine organochlorine pollutants in compost. *Environmental Science and Technology* 31 (10): 2960-2965.
- Strandberg, B.; Bergqvist, P.-A. and Rappe, C. 1998. Dialysis with semipermeable membranes as an efficient lipid removal method in the analysis of bioaccumulative chemicals. *Analytical Chemistry* 70 (3): 526-533.
- Strandberg, B.; Gustafson, P.; Söderström, H.S.; Barregard, L.; Bergqvist, P.-A. and Sallsten, G. 2006. The use of semipermeable determine persistent membrane devices as passive samplers to organic compounds in indoor air. *Journal of Environmental Monitoring* 8 (2): 257-262.
- Stuer-Lauridsen, F. and Dahl, B. 1995. Source of organotin at a marine water/sediment interface - A field study. *Chemosphere* 30 (5): 831-845.
- Stuer-Lauridsen, F. and Kjølholt, J. 2000. Identification of selected hydrophobic organic contaminants in wastewater with semipermeable membrane devices (SPMDS). *Water Research* 34 (13): 3478-3482.
- Stuer-Lauridsen, F. 2005. Review of passive accumulation devices for monitoring organic micropollutants in the aquatic environment. *Environmental Pollution* 136 (3): 503-524.
- Sundberg, H.; Tjärnlund, U.; Åkerman, G.; Blomberg, M.; Ishaq, R.; Grunder, K.; Hammar, T.; Broman, D. and Balk, L. 2005. The distribution and relative toxic potential of organic chemicals in a PCB contaminated bay. *Marine Pollution Bulletin* 50 (2): 195-207.
- Sztamberek-Gola, I.; Grochowski, A. and Chrząszcz, R. 2003. Monitoring of PCDDs, PCDFs, and PAHs in wastewater with use of semipermeable membrane devices (SPMD). *Organohalogen Compounds* 60: 45-48.
- Ter Laak, T.L.; Agbo, S.O.; Barendregt, A. and Hermens, J.L.M. 2006. Freely dissolved concentrations of PAHs in soil pore water: Measurements via solid-phase extraction and consequences for soil tests. *Environmental Science and Technology* 40 (4): 1307-1313.
- Ter Laak, T.L.; Barendregt, A. and Hermens, J.L.M. 2006. Freely dissolved pore water concentrations and sorption coefficients of PAHs in spiked aged, and field-contaminated soils. *Environmental Science and Technology* 40 (7): 2184-2190.
- Tusseau-Vuillemin, M.-H.; Gourlay, C.; Lorgeoux, C.; Mouchel, J.-M.; Buzier, R.; Gilbin, R.; Seidel, J.-L. and Elbaz-Poulichet, F. 2007. Dissolved and bioavailable contaminants in the Seine river basin. *Science of the Total Environment* 357 (1-3): 244-256.
- Utvik, T.I.R.; Durell, G.S. and Johnsen, S. 1999. Determining produced water originating polycyclic aromatic hydrocarbons in North Sea waters: Comparison of sampling techniques. *Marine Pollution Bulletin* 38 (11): 977-989.
- Utvik, T.I.R. and Johnsen, S. 1999. Bioavailability of polycyclic aromatic hydrocarbons in the North Sea. *Environmental Science and Technology* 33 (12): 1963-1969.
- van der Oost, R.; Beyer, J. and Vermeulen, N.P.E. 2003. Fish bioaccumulation and biomarkers in environmental risk assessment: A review. *Environmental Toxicology and Pharmacology* 13 (2): 57-149.
- van der Oost, R.; Verweij, F.; Satumalay, K.; van der Molen, N. and Booij, K. 2004. Fish biomarker responses for the assessment of environmental risks compared to SPMD-bioassay alternatives: Abstract. *Marine Environmental Research* 58 (2-5): 306-307.
- van Drooge, B.L.; Grimalt, J.O.; Booij, K.; Camarero, L. and Catalan, J. 2005. Passive sampling of atmospheric organochlorine compounds by SPMDs in a remote high mountain area. *Atmospheric Environment* 39 (28): 5195-5204.
- van Stee, L.L.P.; Leonards, P.E.G.; van Loon, W.M.G.M.; Hendriks, A.J.; Maas, J.L.; Struijs, J. and Brinkman, U.A.T. 2002. Use of semi-permeable membrane devices and solid-phase extraction for the wide-range screening of microcontaminants in surface water by GC-AED/MS. *Water Research* 36 (18): 4455-4470.
- van Zwieten, L.; Ahmad, N.; Marolt, R.S.; Ayres, M. and Morris, S. 2001. Determination of DDT and its

- metabolites in cattle dip soil available in aqueous phase after remediation. *Journal of Environmental Science and Health Part B-Pesticides Food Contaminants and Agricultural Wastes* 36 (5): 501-516.
- Verweij, F.; Booij, K.; Satumalay, K.; van der Molen, N. and van der Oost, R. 2004. Assessment of bioavailable PAH, PCB and OCP concentrations in water, using semipermeable membrane devices (SPMDs), sediments and caged carp. *Chemosphere* 54 (11): 1675-1689.
- Villeneuve, D.L.; Crunkilton, R.L. and DeVita, W.M. 1997. Aryl hydrocarbon receptor-mediated toxic potency of dissolved lipophilic organic contaminants collected from Lincoln Creek, Milwaukee, Wisconsin, USA, to PLHC-1 (*Poeciliopsis lucida*) fish hepatoma cells. *Environmental Toxicology and Chemistry* 16 (5): 977-984.
- Villeneuve, J.Y.; Niimi, A.J. and Metcalfe, C.D. 1999. Distribution and bioaccumulation of chlorinated diphenyl ethers in a contaminated embayment of Lake Ontario. *Journal of Great Lakes Research* 25 (4): 760-771.
- Villeneuve, D.L.; Curtis, L.R.; Jenkins, J.J.; Warner, K.E.; Tilton, F.; Kent, M.L.; Watral, V.G.; Cunningham, M.E.; Markle, D.F.; Sethajintanin, D.; Krissanakriangkrai, O.; Johnson, E.R.; Grove, R.A. and Anderson, K.A. 2005. Environmental stresses and skeletal deformities in fish from the Willamette River, Oregon. *Environmental Science and Technology* 39 (10): 3495-3506.
- Vinturella, A.E.; Burgess, R.M.; Coull, B.A.; Thompson, K.M. and Shine, J.P. 2004. Use of passive samplers to mimic uptake of polycyclic aromatic hydrocarbons by benthic polychaetes. *Environmental Science and Technology* 38 (4): 1154-1160.
- Voie, Ø.A.; Johnsen, A. and Rossland, H.K. 2002. Why biota still accumulate high levels of PCB after removal of PCB contaminated sediments in a Norwegian fjord. *Chemosphere* 46 (9-10): 1367-1372.
- Vrana, B.; Paschke, A. and Popp, P. 2001. Polyaromatic hydrocarbon concentrations and patterns in sediments and surface water of the Mansfeld region, Saxony-Anhalt, Germany. *Journal of Environmental Monitoring* 3 (6): 602-609.
- Vrana, B.; Paschke, A.; Popp, P. and Schüürmann, G. 2001. Use of semipermeable membrane devices (SPMDs) - Determination of bioavailable, organic, waterborne contaminants in the industrial region of Bitterfeld, Saxony-Anhalt, Germany. *Environmental Science and Pollution Research* 8 (1): 27-34.
- Vrana, B. and Schüürmann, G. 2002. Calibrating the uptake kinetics of semipermeable membrane devices in water: Impact of hydrodynamics. *Environmental Science and Technology* 36 (2): 290-296.
- Vrana, B.; Mills, G.A.; Allan, I.J.; Dominiak, E.; Svensson, K.; Knutsson, J.; Morrison, G. and Greenwood, R. 2005. Passive sampling techniques for monitoring pollutants in water. *Trac-Trends in Analytical Chemistry* 24 (10): 845-868.
- Vrana, B.; Paschke, H.; Paschke, A.; Popp, P. and Schüürmann, G. 2005. Performance of semipermeable membrane devices for sampling of organic contaminants in groundwater. *Journal of Environmental Monitoring* 7 (5): 500-508.
- Waite, D.T.; Droneck, N.R.; Tuduri, L.; Sproull, J.F.; Chau, D.F. and Quiring, D.V. 2005. Comparison of active versus passive atmospheric samplers for some current-use pesticides. *Bulletin of Environmental Contamination and Toxicology* 74 (6): 1011-1018.
- Wang, C.; Wang, Y.; Kiefer, F.; Yediler, A.; Wang, Z. and Kettrup, A. 2003. Ecotoxicological and chemical characterization of selected treatment process effluents of municipal sewage treatment plant. *Ecotoxicology and Environmental Safety* 56 (2): 211-217.
- Wang, J., Bi, Y., Pfister, G., Henkelmann, B., Zhu, K. and Schramm, K-W. (2009) Determination of PAH, PCB, and OCP in water from the Three Gorges Reservoir accumulated by semipermeable membrane devices (SPMD). *Chemosphere*, 75 (8), 1119-1127.
- Wang, Y.; Wang, C. and Wang, Z. 1998. Uptake of moderately hydrophobic chlorophenols from water by semipermeable membrane devices (SPMDs) and by goldfish (*Crassius auratus*). *Chemosphere* 37 (2): 327-339.
- Wang, Y.; Wang, Z.; Liu, J.; Ma, M. and Belzile, N. 1999. Monitoring priority pollutants in the Yangtze River by dichloromethane extraction and semipermeable membrane device (SPMD). *Chemosphere* 39 (1): 113-131.
- Wang, Y.; Wang, Z.; Wang, C. and Wang, W. 1999. Uptake of weakly hydrophobic nitroaromatics from water by semipermeable membrane devices (SPMDs) and by goldfish (*Carassius auratus*).

Chemosphere 38 (1): 51-66.

- Wang, Y.; Wang, Z.; Liu, J.; Ma, M. and Wang, C. 2000. Monitoring toxic organic pollutants in Huaihe River waters using SPE and triolein-SPMD. *China Environmental Science* 20 (2): [vp].
- Wang, Y.; Wang, Z.; Ma, M.; Wang, C. and Mo, Z. 2001. Monitoring priority pollutants in a sewage treatment process by dichloromethane extraction and triolein-semipermeable membrane device (SPMD). *Chemosphere* 43 (3): 339-346.
- Wang, Y.; Huang, Y.; Huckins, J.N. and Petty, J.D. 2004. Compound-specific carbon and hydrogen isotope analysis of sub-parts per billion level waterborne petroleum hydrocarbons. *Environmental Science and Technology* 38 (13): 3689-3697.
- Wang, Z.; Wang, Y. and Ma, M. 2001. Assessing the ecological risk of sediment-associated polychlorinated biphenyls in Huaihe River. *China Environmental Science* 21 (3): 262-265.
- Wang, Z.; Lu, Y.; Wang, Y. and Ma, M. 2002. Assessing the ecological risk of substituted benzenes in Huaihe River, China. *Acta Scientiae Circumstantiae/Huanjing Kexue Xuebao. Beijing* 22 (3): 300-304.
- Wang, Z. and Wang, Y. 2002. Bioaccumulation and biodegradation of chlorophenols in semi-permeable membrane biomimetic sampling device and in Golden Fish (*Carassius auratus*). *Journal of Lake Sciences* 14 (1): 71-76.
- Wang, Z. and Wang, Y. 2002. Simulation of the accumulation and degradation of nitroaromatics in fish by a biomimetic sampling technology. *Acta Hydrobiologica Sinica/Shuisheng Shengwu Xuebao* 26 (3): 209-214.
- Wang, Z.; Wang, Y.; Ma, M.; Lu, Y. and Huckins, J.N. 2002. Use of triolein-semipermeable membrane devices to assess the bioconcentration and sediment sorption of hydrophobic organic contaminants in the Huaihe River, China. *Environmental Toxicology and Chemistry* 21 (11): 2378-2384.
- Wells, J.B. and Lanno, R.P. 2001. Passive sampling devices (PSDs) as biological surrogates for estimating the bioavailability of organic chemicals in soil. *Environmental Toxicology and Risk Assessment: Science, Policy, and Standardization Implications for Environmental Decisions*, at Toronto, Ont. (Canada). ASTM Special Technical Publication. No. 1403.
- Wenzel, K.D.; Vrana, B.; Hubert, A. and Schürmann, G. 2004. Dialysis of persistent organic pollutants and polycyclic aromatic hydrocarbons from semipermeable membranes. A procedure using an accelerated solvent extraction device. *Analytical Chemistry* 76 (18): 5503-5509.
- Werner, D.; Ghosh, U. and Luthy, R.G. 2006. Modeling polychlorinated biphenyl mass transfer after amendment of contaminated sediment with activated carbon. *Environmental Science and Technology* 40 (13): 4211-4218.
- Whyte, J.J.; Karrow, N.A.; Boermans, H.J.; Dixon, D.G. and Bols, N.C. 2000. Combined methodologies for measuring exposure of rainbow trout (*Oncorhynchus mykiss*) to polycyclic aromatic hydrocarbons (PAHs) in creosote contaminated microcosms. *Polycyclic Aromatic Compounds* 18 (1): 71-98.
- Williamson, K.S.; Petty, J.D.; Huckins, J.N.; Lebo, J.A. and Kaiser, E.M. 2002. HPLC-PFD determination of priority pollutant PAHs in water, sediment, and semipermeable membrane devices. *Chemosphere* 49 (7): 703-715.
- Williamson, K.S.; Petty, J.D.; Huckins, J.N.; Lebo, J.A. and Kaiser, E.M. 2002. Sequestration of priority pollutant PAHs from sediment pore water employing semipermeable membrane devices. *Chemosphere* 49 (7): 717-729.
- Xu, J.; Wang, P.; Zhong, X.; Dai, S.U.; Sun, H. and Hao, Y. 2006. Comparison of PAH and nonylphenol uptake by carp (*Cyprinus carpio*) and semipermeable membrane devices (SPMDs) from water. *Bulletin of Environmental Contamination and Toxicology* 77 (2): 211-218.
- Yusa, V.; Pastor, A. and de la Guardia, M. 2005. Microwave-assisted extraction of OCPs, PCBs and PAHs concentrated by semi-permeable membrane devices (SPMDs). *Analytica Chimica Acta* 540 (2): 355-366.
- Yusà, V.; Pastor, A. and de la Guardia, M. 2006. Microwave-assisted extraction of polybrominated diphenyl ethers and polychlorinated naphthalenes concentrated on semipermeable membrane devices. *Analytica Chimica Acta* 565 (1): 103-111.
- Zabiegala, B.; Kot, A. and Namieśnik, J. 2000. Long-term monitoring of organic pollutants in water - Application of passive dosimetry. *Chemia Analityczna* 45 (5): 645-657.
- Zajicek, J.L.; Tillitt, D.E.; Huckins, J.N.; Petty, J.D.; Potts, M.E. and Nardone, D.A. 1996. Application of

- enzyme-linked immunosorbent assay for measurement of polychlorinated biphenyls from hydrophobic solutions - Extracts of fish and dialysates of semipermeable membrane devices. In: *Environmental Immunochemical Methods, ACS Symposium Series*. Vol 646, pp. 307-325.
- Zhang, Y.; Lam, M.H.W.; Fai, P.K.; Richardson, B.J.; Hong, H.-s.; Zheng, G.J. and Wu, R.S.S. 2000. Introduction of a new device for sampling organic pollutants. *Chinese Journal of Analytical Chemistry* 28 (11): 1434-1438.
- Zhang, J. and Zhang, Y. 2001. Development and application of SPMD technology in environmental research. *Marine environmental science/Haiyang Huanjing Kexue. Dalian* 20 (4): 67-73.
- Zheng, J.; O. S.; Pan, L. and Richardson, B.J. 1999. Semi-permeable membrane devices (SPMD) study for marine organic chloro-pesticide pollution monitoring. *Marine environmental science/Haiyang Huanjing Kexue. Dalian* 18 (1): 19-23.
- Zhu, Y.-x.; Zhang, Y.; Zhang, J.; Zhuang, Y.-t.; Poon, K.-F.; Lam, M.H.W.; Hong, H.-s. and Wu, R.S.S. 2001. Monitoring polynuclear aromatic hydrocarbons in sediment porewater by SPMD. *Chinese Journal of Oceanology and Limnology* 19 (4): 382-388.
- Zimmerman, L.R.; Thurman, E.M. and Bastian, K.C. 2000. Detection of persistent organic pollutants in the Mississippi Delta using semipermeable membrane devices. *Science of the Total Environment* 248 (2-3): 169-179.
- Zimmerman, J.R.; Ghosh, U.; Millward, R.N.; Bridges, T.S. and Luthy, R.G. 2004. Addition of carbon sorbents to reduce PCB and PAH bioavailability in marine sediments: Physicochemical tests. *Environmental Science and Technology* 38 (20): 5458-5464.
- Zimmerman, J.R.; Werner, D.; Ghosh, U.; Millward, R.N.; Bridges, T.S. and Luthy, R.G. 2005. Effects of dose and particle size on activated carbon treatment to sequester polychlorinated biphenyls and polycyclic aromatic hydrocarbons in marine sediments. *Environmental Toxicology and Chemistry* 24 (7): 1594-1601.

The following are Letters to the Editor of peer-reviewed journals. The letters discuss aspects of papers that had recently appeared in that particular journal.

n=7

- Lebo, J.A.; Huckins, J.N. and Meadows, J.C. 1996. Critique of "Passive diffusion through polymeric membranes: A novel cleanup procedure for analysis of azinphos-methyl and azinphos-ethyl residues in fruits and vegetables" Ahmad, N. et al. (1995) J. AOAC Int. 78, 1450-1454. *Journal of AOAC International* 79 (4): A78.
- Ahmad, N.; Mandarakas, P.; Guo, L.; Appleby, S. and Bugeno, G. 1996. The authors' response: to "Critique of Passive diffusion through polymeric membranes: A novel cleanup procedure for analysis of azinphos-methyl and azinphos-ethyl residues in fruits and vegetables" Ahmad, N. et al. (1995) J. AOAC Int. 78, 1450-1454". *Journal of AOAC International* 79 (4): A78.
- Huckins, J.N.; Petty, J.D.; Lebo, J.A.; Orazio, C.E. and Prest, H.F. 1997. Comment on "Accumulation of organochlorine pesticides and PCBs by semipermeable membrane devices and *Mytilus edulis* in New Bedford Harbor". *Environmental Science and Technology* 31 (12): 3732-3733.
- Shea, D. and Hofelt, C.S. 1997. Response to "Comment on Accumulation of organochlorine pesticides and PCBs by semipermeable membrane devices and *Mytilus edulis* in New Bedford Harbor". *Environmental Science and Technology* 31 (12): 3734-3735.
- Prest, H.F.; Petty, J.D. and Huckins, J.N. 1998. Validity of using semipermeable membrane devices for determining aqueous concentrations of freely dissolved PAHs: Letter to Editor. *Environmental Toxicology and Chemistry* 17 (4): 535-536.
- Gustafson, K.E. and Dickhut, R.M. 1998. The authors' response: to "Validity of using semipermeable membrane devices for determining aqueous concentrations of freely dissolved PAHs". *Environmental Toxicology and Chemistry* 17 (4): 536.
- Schwartz, R.M. 2006. Comparison of mussels and semi-permeable membrane devices as inter-tidal monitors of polycyclic aromatic hydrocarbons at oil spill sites (Boehm et al. 2005) letter. *Marine Pollution Bulletin* 52 (1): 121-121.